

Measurement of Jet Inclusive Cross Section using the KT Algorithm in CDF RUN2






R. Lefevre, M. Martinez, O. Norniell
IFAE

Introduction

- Context
 - $0.1 < |Y| < 0.7$
 - $D = 0.5, 0.7$ and 1.0
- Plan
 - Used data
 - Cuts
 - Trigger study
 - P_T corrections
 - Unfolding
 - Systematics
 - Results

Used data


- Datasets xxxx0c
 - ST5, J20, J50, J70 and J100 Triggers
- Runs used
 - V4 of QCD good run list
 - Excluded runs
 - Processing problem (why?)
 - [154594, 155895] and [156099, 160151]
 - Vz proble in production software
 - ≥ 166328 : loose around 35pb^{-1}
 - Only for ST5: unconstant L2 prescale
 - < 147870
- Integrated luminosity: 145pb^{-1} (129pb^{-1} for ST5)




Cuts

- $0.1 < |Y| < 0.7$
- $|V_z| < 60\text{cm}$
 - With at least 3 associated tracks
- Missing E_T significance
 - Cut function of P_T


D=0.5 PYTHIA: vertex cut efficiencies



PYTHIA: missing E_T significance cut study




PYTHIA: missing E_T significance cut efficiency




Trigger study

- Study done for L1, L2 and L3
 - Use high PT muon trigger to get ST5 efficiency
- Minimal raw P_T used


| Trigger | D = 0.5 / 0.7 / 1.0 |
|---------|---------------------|
| ST5 | 23 / 27 / 31 |
| J20 | 27 / 35 / 41 |
| J50 | 62 / 62 / 72 |
| J70 | 83 / 96 / 96 |
| J100 | 127 / 127 / 146 |

ST5(L1) / High P_T muon


J15(L2) / ST5




J20(L3) / ST5




ST10(L1) / J20




J40(L2) / J20




J50(L3) / J20




J60(L2) / J50




J70(L3) / J50




J90(L2) / J70







J100(L3) / J70








D=0.5: Raw measurement




















P_T corrections

- From PYTHIA:
 - Matching Calorimeter/Hadron Jets
 - $\Delta R < D$ in ϕ -Y plan
 - $\langle P_T \text{ Raw} - P_T \text{ Had} \rangle$ vs $\langle P_T \text{ Raw} \rangle$
- Pile-up correction
 - $\Psi \times$ Number of additional primary vertices
 - Ψ values extract from dedicated studies performed with cone algorithm
 - 100% systematics uncertainty
 - 0.400, 0.781 and 1.581 GeV/c respectively for $D = 0.5, 0.7$ and 1.0












D=0.5 PYTHIA: P_T correction




D=0.5 PYTHIA: result of P_T correction




D=0.5 PYTHIA: P_T correction




D=0.5 PYTHIA: result of P_T correction




D=0.5: before pile-up correction




D=0.7: before pile-up correction




D=1.0: before pile-up correction




D=0.5: after pile-up correction



D=0.7: after pile-up correction




D=1.0: after pile-up correction



Unfolding

- A simple bin by bin correction is used
 - $N_{\text{hadron level}} / N_{\text{calorimeter level (corrected } P_T)}$
- Extract from PYTHIA


Unfolding obtained from PYTHIA




Systematics

- $|V_z|$ cut
 - ± 5 cm: effect lower than 1%
- Number of tracks associated to primary vertex
 - ± 1 tracks: effect of the order of 0.1%
- Missing E_T significance cut
 - $\pm 10\%$ on missing E_T scale
 - ⊗ $\pm 3\%$ on jet energy scale: effect lower than 0.3%
- MC statistics: lower than 2%
- $\pm 3\%$ on jet energy scale
- $\pm 100\%$ on pile-up correction
- $\pm 3\%$ on lower edge of each bin
- Unfolding: PYTHIA vs HERWIG


D=0.5: Jet Energy Scale changed of +/- 3%




Jet Energy Scale changed of +/- 3%




Pile-up energy correction changed of +/- 100%




D=0.5: Lower edge of each bin moved of +/- 3%



D=0.5: Lower edge of each bin moved of +/- 3%




Unfolding systematic from Herwig and Pythia comparison




Results


CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY




CDF RUN2 PRELIMINARY



CDF RUN2 PRELIMINARY



CDF RUN2 PRELIMINARY



Comments

- Systematic from MC simulation of the calorimeters still missing
- ST5 has lower integrated luminosity
 - To be specified on plots?
 - Simply removed those points?
- In progress
 - NLO comparisons